

## Araxa carbonatite deposit and its lateritization

Masaharu KAMITANI\* and Hideo HIRANO\*\*

KAMITANI, M. and HIRANO, H. (1990) Araxa carbonatite deposit and its lateritization. *Bull. Geol. Surv. Japan*, vol. 41 (11), p. 595-604.

**Abstract:** Araxa niobium-bearing carbonatite deposit occurring in Barreiro carbonatite was studied preliminarily. The Barreiro (Araxa) complex located at approximately 6 kilometers south of Araxa city, Minas Gerais, Brazil, is intruded into the Araxa group of the lower Proterozoic system and is composed mainly of beforosite, phoscolite, sovite, glimmerite and fenite. This complex including the fenite zone with about 5.5 kilometers in diameter and both of the complex and the host Proterozoic systems have been deeply weathered and consequently the fresh rocks are scarcely observed in this area. According to exploration drilling data in the mining site, the thickness of the lateritized mantle vary from a several to two hundred meters.

The lateritization has brought out the enrichment of niobium, titanium and phosphorus. This study reveals the concentration ratios of the ore components based on the chemical compositions between the primary carbonatite and the weathered mantle in the Barreiro complex.  $ZrO_2$  in the beforosite has increased about 8 times in the weathered mantle and  $SiO_2$ ,  $Al_2O_3$  and BaO are 6 to 9 times. It is therefore concluded that the formation of anomalously high-grade niobium ore deposit was clearly caused by the strong lateritization.

### 1. Introduction

Carbonatites that have been given special attention not only to the petrological interests but also to the economical importance are distributed interior and marginal portions of continental areas. There are more than five hundreds localities of carbonatites in the world (KAMITANI and HIRANO, 1985). The unique mineral resources of carbonatites have high potentiality for niobium, rare-earth, titanium, zirconium and phosphorus (DEANS, 1978). Especially, niobium production from carbonatite ore deposits shares more than 98 per cent of the world production. Araxa and Catalao mines are famous for the niobium-bearing carbonatite ore deposits, and the other Brazilian carbonatite complexes have been paid attention to their mineral resources of phosphorus, titanium and rare-earth.

In the southern Brazil, many carbonatite-alkaline/alkaline complexes are recognizable particularly in the peripheries of the Parana basin (SCHOBENHAUS, 1981) which consists mainly of Paleozoic to Mesozoic sediments with tholeiitic basalts (Fig. 1). The tholeiitic basalts were formed in Triassic to Jurassic age and the many carbonatite-alkaline/alkaline complexes were intruded principally in Jurassic to Cretaceous age (ULBRICH and GOMES, 1981). The Parana basalts are believed to be a result of large-scale fissure eruptions related to the break up of the Gondwana old continent. In the Angola and the South Africa similar intrusion age of carbonatite complexes are distributed (MARSH, 1973; ULBRICH and GOMES, 1981). Most of those K-Ar ages in the southern Brazil ranges from 135 to 60 Ma (AMARAL *et al.*, 1967). HERZ (1977) states that the northwest alignment of the complexes in the Cretaceous age including the Araxa and Catalao carbonatites are resulted from one of the failed arms of the triple junctions. It might be kind of marginal fracture zones that is roughly across at right angle to the plate margin

\* Geological Museum, Geological Survey of Japan.

\*\* Mineral Resources Dept., Geological Survey of Japan.




Fig. 1 Geologic map of southern Brazil.

(FRANCHETEAU and PICHON, 1972).

The carbonatite-alkaline/aklaline complexes in Brazil were grouped into eight types by ULBRICH and GOMES (1981). All of the carbonatite-bearing complexes belonging to the Type III of Ulbrich and Gomes are always accompanied with glimmerites, dunites, peridotites and pyroxenites.

The Barreiro carbonatite complex is accompanied with niobium-bearing ore body in the central part of the complex, as well as phosphorus and rare-earth ore deposits in the north to the northeast (BERBERT, 1984). In this paper the secondary enrichment of niobium was preliminarily studied and the concentration process is discussed based on the chemical analysis data.

## 2. Geologic setting

### 2.1 Regional geology

Barreiro (Araxa) carbonatite complex is located at about 300 kilometers west of Belo Horizonte, the third largest city of the Brazil. The mining site of the Araxa niobium ore deposit has been constructed just south of Araxa city on the high-land area with the altitude from 1,000 to 1,200 meters.

There are five carbonatite alkaline complexes in the vicinity of Araxa city; Salitre I, II, Serra Negra, Barreiro and Tapira from north to south (Fig. 2). Their K-Ar ages according to HASUI and CORDANI (1968) indicate late Cretaceous but vary from 70 to 91 Ma.

The Salitre I, II, and Serra Negra complexes intruded into the Paraopeba formation of upper Proterozoic system are located at approximately 70 kilometers north of the Araxa. Those complexes with about 10 kilometers in diameter have been completely lateritized and therefore

no fresh rock is recognizable in and around the complexes. Results of exploration trenches and drills indicate that the weathered mantle on the complexes contains an intermediate to high-grade of titanium ore deposits (Table 1).

The Tapira complex is located at about 30 kilometers southeast of the Araxa and intruded into the Canastra formation of middle Proterozoic system. This complex with about 7 kilometers in diameter is presently under-developing by open pit as a titanium mine, but it has also considerable amount of niobium and phosphorus associated with the titanium deposit.

The geology of this area including the five carbonatite complexes consists mainly of Archean undifferentiated system, middle Proterozoic system of Araxa, Canastra and Ibia formations, and Paraopeba formation of upper Proterozoic. These crystalline basements are unconformably covered with the Serra Jeral formation of late Jurassic in the west and the Paulu formation in the east (DNPM, 1978).

The Serra Jeral formation is characterized by a thick pile of tholeiitic basalts and their pyroclastics. This volcanism started with fissure eruption about 147 Ma ago in the Santa Catarina area and its peak is thought to be between 120 and 130 Ma (CORDANI and VANDROS, 1967). In the late stage of the volcanism many diabasic dykes were intruded into the basement along NW trending fractures which are parallel to that of magnetic linearments (SCHOBENHAUS, 1981). The NW trending faults are also observable around the Salitre complexes, but obscure around the Barreiro and Tapira.

### 2.2 Geology and occurrence of the Araxa niobium ore deposit in the Barreiro complex

Table 1 Mineral resources of carbonatite complexes in Araxa area

	Niobium		Titanium		Phosphorus		Rare earth	
	$\times 10^6$ t,	%Nb <sub>2</sub> O <sub>5</sub>	$\times 10^6$ t,	%TiO <sub>2</sub>	$\times 10^6$ t,	%P <sub>2</sub> O <sub>5</sub>	$\times 10^6$ t,	%RE <sub>2</sub> O <sub>3</sub>
ARAXA	462	2.48			460	15.07	560	10.5
TAPIRA	165	1.76	325	16.3	991	9.04		
SALITRE I			84	23.3				
SALITRE II			92	13.0				




Fig. 2 Geologic map of Araxa area, Minas Gerais.

The Araxa mine area is underlain by Precambrian undifferentiated system and lower Proterozoic Araxa group. The former consisting of gneisses, gneissose granites, migmatites, quartzites, amphibolites and schists, is distributed as a basement in the northern to northeastern part of this area. The Araxa group comprising of mica

schists, quartzites and amphibolites is widely distributed in the area and overlies unconformably the Precambrian undifferentiated system. Geologic structure of this group around the Barreiro carbonatite complex has been deformed by the intrusion of the complex and then the surrounding quartzite and mica schist




Fig. 3 Geologic map of Barreiro (Araxa) carbonatite complex. Nb, P and REE ore deposits developed in the weathered mantle are projected in the map.

beds were turned up along the wall of the intrusive body. A ring-like structure around the complex is clearly recognizable (Fig. 3).

The Barreiro carbonatite shows a concentric zonation; carbonatites concentrate in the core zone and glimmerite in the outer zone. The core zone of the complex consists mainly of befor-site, phoscolite, glimmerite and sövite. Fenitization which is resulted from alkali metasomatism of the intrusive body is observable as a circular halo in the wall rock as seen in Fig. 3.

Glimmerite of which main constituent minerals are phlogopite and dolomite might be an alteration product (GITTINS *et al.*, 1975) from ultramafic rocks formed by a kind of alkali metasomatism due to carbonatite magma intrusion. Small amount of magnetite and apatite are contained within the glimmerite but pyrochlore is very little. Clinopyroxene is reported from some deep drilling cores as relicts replaced by phlogopite. No olivine has been observed but serpentinite occurs as a relict in some drilling

cores (FILHO *et al.*, 1984).

Beforsite which is dolomitic carbonatite contains mainly dolomite and with variable amount of calcite and ankerite. Apatite, pyrochlore, phlogopite and alkali-amphiboles are associated within the rock. According to FILHO *et al.* (1984), phoscolite-like rocks are recognizable in the befor-site zones within the core of the complex as brecciated masses containing apatite, phlogopite, magnetite and pyrochlore, but olivine and pyroxene minerals seen in Phalaborwa and Sokli carbonatite complexes (Palabora Mining Company, 1976; VARTIAINEN and PAARMA, 1979) are not observed. In the Barreiro complex the phoscolite associated with pyrochlore is always cut by carbonatite veins.

Sövite corresponding to calcite carbonatite is observed in the outer zone of the complex. This rock contains a little amount of apatite, pyrite and pyrrhotite, and is often intruded into the fenitized wall rock as small dykes and veins.

In the case of the Barreiro complex, detailed

occurrence and relationship between the rocks in each stage of the carbonatite magmatism have not been clear because of perfect decomposition of the complex by the chemical weathering. In general, fractional crystallization process of carbonatite magma yields sövite, alvikite and beforsite from early to late stage (LEBAS, 1981).

As stated above, the crystallization sequence of the Barreiro complex itself might be estimated as follows; calcite carbonatite and phoscorite, beforsite, and then ankeritic carbonatites.

### 2.3 Ore minerals

Main niobium minerals of the Barreiro carbonatite complex are pyrochlore and bariopyrochlore. The former occurs mainly in the inner core zone consisting of beforsite, and phoscorite. The latter mineral, on the other hand, is thought to be formed in the lateritized mantle of the complex. Although soil samples from the lateritized mantle were collected from the open pit, no fresh carbonatite was obtained from the surface by one of the authors (M.K.). Therefore detailed description on the occurrence and character of the original ore minerals can not be described here.

As seen in Table 2, the main constituent minerals of the lateritized mantle are barite, goethite, hematite, magnetite, anatase, apatite, bariopyrochlore, gibbsite, alunite and kaolinite in order of abundance. A small amount of silica mineral is observed in some part of the mantle. Calcite, dolomite, ankerite and phlogopite which are the main constituent minerals of the fresh carbonatitic rocks were not observed in the weathered mantle. Monazite, magnetite and zircon/baddeleyite are observable as minor constituent minerals in both the carbonatite and lateritized mantle, since these minerals have a resistance to the chemical weathering.

### 3. Migration of chemical composition and secondary enrichment in the niobium ore deposit

Lateritic weathering mantle have been distinctly developed in the whole area of the Brazil. Araxa and its surrounding are also subjected to the strong lateritization and then no

Table 2 Minerals in Araxa niobium ore deposit

Profile	Depth	Mineral Composition
Reddish laterite (overburden)	0-20 m	goethite, hematite, magnetite, anatase, gibbsite, vermicurite, alunite, kaolinite, silica minerals
Laterite (ore body)	20-100 m	goethite, hematite, magnetite, barite, apatite, anatase, bariopyrochlore
Beforsite & phoscolite Original rock (carbonatite)	>100 m	dolomite, calcite, ankerite, magnetite, apatite, perowskite, phlogopite, pyrite, pyrrhotite, pyrochlore

hard fresh rock is seen in and around the mine area except for river terraces. The thickness of the weathered mantle is estimated about 100 meters in average based on the result of many exploration drillings by CBMM (GROSSI and TORRES, 1978). In some part within the Barreiro complex, the weathered mantle is over 200 meters in thickness (FILHO *et al.*, 1984).

It is well known phenomenon that laterite-type of nickel and aluminum deposits in the tropical areas are formed by secondary enrichment of strong chemical weathering. Some high-grade of ore deposits containing niobium, titanium and phosphorus have been also considered as a result of the chemical weathering for the enrichment.

The Araxa niobium deposit in the Barreiro complex is famous for the highest-grade and the biggest ore reserves in the world (PERRAULT and MANKER, 1981; BERBERT, 1984). The lateritization is considered to have played an important role for the secondary enrichment on niobium and others.

SILVA *et al.* (1979) reported a chemical composition of the representative carbonatite from the beforsite zone, which is given in Table 3. Five lateritic soil samples from the niobium-bearing ore horizon and two reddish lateritic soil samples from the overburden were collected by one of the authors and chemically analysed. Average chemical compositions of each lateritic horizon are presented in Table 3.

It is clear that MgO, CaO, CO<sub>2</sub> comprising dolomite and calcite which are the main consti-

Table 3 Chemical compositions of lateritized mantle and carbonatite in Araxa niobium deposit

	Carbonatite <sup>1)</sup> Beforsite & Phoscolite	Laterite <sup>2)</sup> (Ore body)	Reddish <sup>3)</sup> laterite (overburden)
SiO <sub>2</sub>	1.40%	2.50%	16.55%
Al <sub>2</sub> O <sub>3</sub>	1.80	1.23	29.97
Fe <sub>2</sub> O <sub>3</sub>	8.14	40.70	32.52
FeO	17.60	—	—
MnO	1.08	0.79	0.11
MgO	14.50	0.27	0.04
CaO	9.20	0.49	0.62
Na <sub>2</sub> O	0.14	0.30	0.31
K <sub>2</sub> O	0.42	0.14	0.18
P <sub>2</sub> O <sub>5</sub>	0.69	1.98	1.95
BaO	1.95	26.74**	0.76
SrO	0.59	0.31	0.26
TiO <sub>2</sub>	2.30	2.59	3.72
ZrO <sub>2</sub>	0.02	0.14	0.16
Nb <sub>2</sub> O <sub>5</sub>	0.72	3.00*	0.40*
H <sub>2</sub> O <sup>+</sup>	3.57	3.33	10.04
H <sub>2</sub> O <sup>-</sup>		0.28	0.96
CO <sub>2</sub>	33.00	1.71	1.04
SO <sub>3</sub>	1.95	13.50**	0.34**
Others	0.66		
total	99.73	100.00	100.00

1) SILVA *et al.*, 1989.

2) Average of 2 samples from -10 and -20 m levels.

3) Average of 5 samples from -30 and -40 m, and three from -50 m levels.

\* Estimated value; \*\* Estimated value as BaSO<sub>4</sub>.

tuent minerals of the carbonatite, have been leached out from the weathered mantle and that magnetite components have been completely oxidized. The original volume of the carbonatite must have been remarkably decreased owing to the loss of MgO, CaO and CO<sub>2</sub> from the mantle (Fig. 4). Silicates and some oxides would have been remained in the mantle as secondary minerals although these were once decomposed perfectly by the weathering. That is, a primary mineral assemblage of magnetite, phlogopite, barite, apatite and pyrochlore were converted to goethite, hematite, gibbsite, barite, anatase, bariopyrochlore, quartz and amorphous silica mineral (Table 2).

A distinct volume depletion contributes to the grade-up of Nb<sub>2</sub>O<sub>5</sub> content in the lateritized mantle. This is one of the reasons for the second-




Fig. 4 Vertical change of chemical compositions due to lateritic weathering in Araxa niobium deposit. Data from Table 3.

dary enrichment of the ore deposit. In the case of the Araxa ore deposit, the primary ore grade (0.72% Nb<sub>2</sub>O<sub>5</sub>, SILVA *et al.*, 1979) in the fresh ore body has increased into 2.48 percent Nb<sub>2</sub>O<sub>5</sub> in lateritized mantle (Annex 2 of RODRIGUES and LIMA, 1984). In some part of the present open pit, an ore horizon contains more than 3.0 percent Nb<sub>2</sub>O<sub>5</sub> (personal communication from Mr. Souza, Geology Department, CBMM). SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> and BaO contained in phlogopite and barite are perfectly removed and then some of them are enriched into the mantle. SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> and BaO became approximately 6 to 9 times in weight per cent in the mantle. The original rock volume during the lateritization is estimated to become about 15 percent.

In general, zircon and baddeleyite are the most resistant minerals against the lateritization. Zirconium oxide content of the mantle increases into about 8.0 times compared with that of the fresh carbonatite. Hence, decreasing of the original rock volume is assumed to be more than 87.5 percent. Zirconium mineral is therefore considered as one of the indicative minerals to reveal a degree of the secondary enrichment during the lateritization.

REEDMAN (1984) assessed the secondary enrichment on the case of Sukulu carbonatite complex which was covered by thick residual soil containing pyrochlore, apatite and baddeleyite. He points out that pyrochlore and zirconium oxide in the soil were concentrated about 5 times to that of the fresh rock.

According to Table 4, the concentration ratios of Al<sub>2</sub>O<sub>3</sub>, BaO and ZrO<sub>2</sub> are essentially the same in each other and are 8 times. Total Fe<sub>2</sub>O<sub>3</sub>,

Table 4 Chemical variation under lateritization in Araxa niobium deposit. A, B, and C are calculated as dealkaline and anhydrous basis from the data in Table 3

	Carbonatite	Laterite	Reddish laterite	Whole laterite (W)	
	(beforsite & phoscolite)	(orebody)	(overburden)	$0.6 \times B + 0.4 \times C$ =W	Concentration ratio
	C	B	A	W	W/C
SiO <sub>2</sub>	1.5	2.7	19	9.2	6.1
Al <sub>2</sub> O <sub>3</sub>	1.9	1.3	35	15	7.9
T.Fe <sub>2</sub> O <sub>3</sub>	29	44	37	41	1.4
MnO	1.1	0.84	0.13	0.56	0.51
MgO	15	0.29	0.05	0.19	0.01
CaO	9.6	0.52	0.71	0.60	0.06
P <sub>2</sub> O <sub>5</sub>	0.72	2.1	2.2	2.1	2.9
BaO	2.0	28	0.87	17	8.5
TiO <sub>2</sub>	2.4	2.8	4.3	3.4	1.4
ZrO <sub>2</sub>	0.02	0.15	0.18	0.16	8.0
Nb <sub>2</sub> O <sub>5</sub>	0.75	3.2	0.46	2.1	2.8
CO <sub>2</sub>	34	—	—	—	—
SO <sub>3</sub>	2.0	14	0.39	8.6	4.3
Total	100	100	100	100	—

P<sub>2</sub>O<sub>5</sub>, TiO<sub>2</sub> and Nb<sub>2</sub>O<sub>5</sub> vary from 1.4 to 2.9 times. Some of these components may have been transported to the outside of the system by the ground water, because the minerals of the primary rocks have been mostly decomposed and some of them changed into secondary minerals such as goethite, hematite, apatite, anatase, bariopyrochlore, gibbsite, kaolinite and amorphous silica.

#### 4. Concluding remarks

The Barreiro carbonatite complex is a cylindrical mass with about 5 kilometers in diameter standing in the Proterozoic Araxa group. The complex consisting of beforsite, phoscolite, glimmerite, sövite and fenite intruded into the Proterozoic Araxa group during Cretaceous time which is shown by K-Ar phlogopite age of 80–83 Ma (HASUI and CORDANI, 1968). During the Cenozoic era the complex was strongly weathered and the top portion was converted to thick lateritized mantle with 100 meters thick. Main mineral resources of phosphorus, rare-earth; titanium and niobium originally contained in the complex were remarkably enriched

within the lateritized mantle. Pyrochlore as a niobium mineral, for example, is contained mainly in the phoscolite of the beforsite zone. The Nb<sub>2</sub>O<sub>5</sub> content of the fresh rock has been estimated approximately 0.8 percent based on the CBMM's exploration drillings. Niobium-bearing zone continues about 900 meters beneath the present surface. In the lateritized mantle, on the other hand, niobium mineral is bariopyrochlore which is niobium-bearing bario-hydroxide. Nb<sub>2</sub>O<sub>5</sub> grade in the mantle with about 100 meters thick is about 2.1 percent in average. Therefore, it is concluded that the strong lateritization played a major role for the niobium concentration in the Araxa deposit.

#### References

- AMARAL, G. BUSHEE, J., CORDANI, U. J., KAWASHITA, K. and REYNOLDS, J. H. (1967) Potassium-argon ages of alkaline rocks from southern Brasil, *Geochim. Cosmochim. Acta*, vol. 31, p. 117–142.
- BERBERT, C. O. (1984) Carbonatites and associated mineral deposits in Brazil. In *Proced Centennial Internat. Symp. On*



- Geologic Evolution, Resources and Hazards (ed. Y. SHIMAZAKI), *Special report of Geological Survey of Japan*. no. 263, p. 269-291.
- CORDANI, U. G. and VANDOROS, P. (1967) Basaltic rocks of the Palana basin, in Problems in Brazilian Gondwana Geology. *Univ. Fed. Parana, Inst. Geol.*, p. 207-231.
- DEANS, T. (1978) Mineral production from carbonatite complexes: world review. In proceed. of the 1st Intern. Sympo. on carbonatite (ed. RAMOS J. R. de A.), Pocos de Cardas, M.G., Brazil, p. 123-133.
- DNPM-Departamento Nacional da Producao Mineral (1978) Carta Geologia ao Milionesimo, Belo Horizonte, scale in 1:1,000,000.
- FILHO, A. I., LIMA, P. R. A. dos S. and SOUZA, O. M. (1984) Complexos carbonatiticos do Brasil: Aspectos da geologia do complexo carbonatitico do Barreiro, Araxa, MG, Brasil. p. 19-44. *Companhia Brasileiro de Metalurgia e Mineracao*, Sao Paulo.
- FRANCHETEAU J. and PICHON, X. L. (1972) Marginal fracture zones as structural framework of continental margin in South Atlantic ocean. *Amer. Assoc. Petrol. Geophys. Bull.*, vol. 56, p. 991-1007.
- GITTINS, J., ALLEN, C. R. and COOPER, A. F. (1975) Phlogopitization of pyroxenite; its bearing on the composition of carbonatite magmas. *Geol. Mag.* vol. 112, p. 503-507.
- GROSSI SAD, J. H. and TORRES, N. (1978) Geology and mineral resources of the Barreiro complex, Araxa, Minas Gerais. In Proceed. of the 1st Intern. Symp. on carbonatite (ed. RAMOS J. R. de A.), Pocos de Caldas, M.G., Brazil, p. 307-312.
- HASUI, Y. and CORDANI, U. G. (1968) Idades Potassio-argonio de rochas eruptivas Mesozoicas do oeste mineiro e sul de Goias. In Congress Brasileiro de Geologia, 22, Beol Horizonte, p. 139-143.
- HERZ, N. (1977) Timing of spreading in the South Atlantic: information from Brazilian alkalic rocks. *Geol. Soc. Am. Bull.*, vol. 88, p. 101-112.
- JAGER, E. *et al.* (1959) A hydrated barium-strontium pyrochlore in a biotite rock from Panda Hill, Tanganyka. *Miner. Mag.*, vol. 32, p. 10-25.
- KAMITANI, M. and HIRANO, H. (1985) Carbonatite deposits in Brazil (part II). *Chishitsu News*, no. 372, p. 1-16 (in Japanese).
- LEBAS, M. J. (1981) Carbonatite magmas. *Mineral. Mag.* vol. 44, p. 133-140.
- MANTOVANI, M. S. M., *et al.* (1985) Trace element and strontium isotope constraints on the origin and evolution of Parana continental flood basalts of Santa Catarina State, Southern Brazil. *Jour. Petrology*, vol. 26, part 1, p. 187-209.
- MARSH, J. S. (1973) Relationships between transform directions and alkaline igneous rock linearments in Africa and South America. *Earth and Planet. Sci. Letters*, vol. 18, p. 317-323.
- Palabora Mining Company Limited Mine Geological and Mineralogical Staff (1976) The geology and the economic deposits of copper, iron and vermiculite in the Palabora igneous complex: a brief reviews. *Econ. Geol.* vol. 71, p. 177-192.
- PERRAULT, G. and MANKER, E. A. (1981) Geology and mineralogy of niobium deposits. In Proceed. of the Intern. Symp. of Niobium 1981, San Francisco, U.S.A., 127p.
- REEDMAN, J. H. (1984) Resources of phosphate, niobium, iron and other elements in residual soils over the Sukulu carbonatite complex, Southeastern Uganda. *Econ. Geol.* vol. 79, p. 716-724.
- RODRIGUES, C. S. and LIMA, P. R. A. dos S. (1984) Complexos carbonatiticos do Brasil, p. 1-17. CBMM. Sao Paulo.
- SCHOBENHAUS, C. (1981) Geologic map of Brazil and adjoining ocean floor including mineral deposits. Scale 1:2,500,000. 4 sheets, DNPM, Ministerio das Minas e Energia.
- SILVA, A. B., MARCHETTO, M. and SOUZA, O. M. (1979) Geology of the Araxa (Barreiro) carbonatite. 17p. *CBMM Report*.
- ULBRICH, H. H. G. J. and GOMES, C. B. (1981) Alkaline rocks from continental Brazil. *Earth Sci. Rev.* vol. 17, p. 135-154.
- VARTIAINEN, H. and PAARMA, H. (1979) Geological characteristics of the Sokli carbonatite complex, Finland. *Econ. Geol.* vol. 74, p. 1296-1306.

アラシャカーボナタイト鉱床とラテライト化作用

神谷雅晴・平野英雄

要 旨

アラシャニオブ鉱床(ブラジル国ミナスジェライス州)の風化土壌の鉱物組成と化学組成を予察的に調べ以下の結果を得た。風化殻は、Al, Fe, Si の濃集する赤色ラテライト層とその下位に発達する Ba, Nb, P の濃集するラテライト層に区分され、それぞれに特徴的な風化生成鉱物が形成されている。ニオブ鉱床形成の主因は、ラテライト化作用により、ニオブが特定の風化土壌層に効果的に濃集したためと解釈された。

(受付：1990年3月27日；受理：1990年9月1日)