

Trigger and Mechanism of Co-seismic Groundwater Level Changes in the Togari350 well, central Japan

Yasuhiro Asai

**Tono Research Institute of Earthquake Science,
Association for the Development of Earthquake Prediction**



The 6th Taiwan - Japan Joint Workshop on Hydrological and Geochemical Research for Earthquake Prediction
26-27 September, 2007, Tainan, Taiwan

Contents




- Continuous groundwater level (GWL) observation
- Feature of co-seismic GWL changes
- Relationship between dynamic strain/tilt observation and co-seismic GWL changes
 - Finding of the threshold of dynamic strain/tilt variations
- Applying the 1-dimensional finite porous aquifer model
- Mechanism of the co-seismic GWL change

Togari Crustal Activity Borehole Observatory (TGR350)




Borehole profile, Instruments, and geological and hydrological environment in and around TGR350 and DH-2.



Pressure transducer groundwater level gauge (OYO Corporation product)

Tow Expected common confined aquifers




The multi-component borehole instrument (Ishii et al, 2002)

Groundwater Level : Since May,1998 [1-hour]



Crustal movement (Strain, Tilt): Since January,1999 [10-sec] ; July,2000 [1-Hz]

TGR350 and DH-2, groundwater level (hourly record)

We observed 17 groundwater level changes in response to local and distant earthquakes.




Locations of TGR350 and epicenters




Feature of co-seismic GWL changes

The common feature of **all co-seismic GWL changes** is 'rise'




Tidal component and Atmospheric pressure response are removed by using BAYTAP-G program.

Peak Amp. - Time to Peak




All normalized all co-seismic groundwater level changes



This result suggest that **the source for co-seismic changes has a linear response to the input.**


Relationship between dynamic strain/tilt variation and co-seismic GWL changes

TGR350 Groundwater Level, Tilt and Strain records
2003/09/26 04:50 SE OFF TOKACHI Mj8.0




GWL rise after the passage of the seismic waves (dynamic strain/tilt variations).

TGR350 Groundwater Level, Tilt and Strain records
2004/12/26 9:58 SUMATRA-ANDAMAN Mw9.0




Comparison of Co-seismic GWL change and Dynamic strain/tilt variations

p-to-p Amp.of Strain - Peak Amp.



p-to-p Amp.of Tilt-down - Peak Amp.



Large dynamic strain/tilt variations

Large Co-seismic change

Verification of the threshold

$$M_{JMA} = -0.45 + 2.45 \log_{10} D$$


(Haibara; Matsumoto and Roeloffs, 2003)

$$M_{JMA} = -1.0 + 2.75 \log_{10} D$$


(TGR350; This study)

However, there are many earthquakes caused no co-seismic GWL changes even when magnitude M_{JMA} and D satisfy above the relation.


We check the peak-to-peak amplitudes of **142 dynamic strain/tilt variations** that caused no co-seismic GWL changes (blue mark) in the period July 2000 to December 2004.



Groundwater level changes and peak-to-peak amplitudes of the dynamic strain variations and tilt-down variations in 2004




Discovery of the threshold values of approximately 3×10^{-7} strain and 2×10^{-4} radian.



Geological and hydrological information in and around TGR350.


Modified from JNC (2003).



Steady state flow

NNW fault


Low permeability



Applying the Roeloffs(1998)'s mechanism

---Diffusion of a localized co-seismic pressure increase in an isotropic homogeneous 1-dimensional finite porous aquifer

Comparison of Earthquake responses and Models



Observed (black lines)
Theoretical (red dashed line)


The head, $h_s(x,t;K_h,S_s)$, satisfies the diffusion equation

$$\frac{\partial^2 h_s}{\partial x^2} = \frac{S_s}{K_h} \frac{\partial h_s}{\partial t} = \frac{1}{c_h} \frac{\partial h_s}{\partial t}$$

Horizontal hydraulic diffusivity $Ch=Kh/S_s$ (m^2/sec)


Kh : hydraulic conductivity [m/sec]

S_s specific storage [m^{-1}]




- 
- Observed co-seismic GWL changes.
 - Discovery of the **threshold values of dynamic strain/tilt variations**.
 - **Geological and hydrological information in and around TGR350.**

We propose a **realistic mechanism** of the co-seismic GWL changes.



NNW fault (Low permeability)

TGR350



Conclusions

- During the period from August, 1998 to June 2005, 17 co-seismic groundwater level changes are observed in TGR350, Central Japan.
All changes are 'rise'. The elapsed time of the peak is in proportion to the peak amplitude of Co-seismic GWL changes.
- Peak amplitude of co-seismic groundwater changes are in proportion to the peak-to-peak amplitude of dynamic strain/tilt variations above the certain threshold values.
- We propose the realistic mechanism of Co-seismic groundwater level changes, which is consistent with geological and hydrological information in and around TGR350.